

On the Role of Exchange Interaction in Magnetic Ordering and Conductivity of Manganites

M.V.Krasinkova*

Ioffe Physical Technical Institute, Russian Academy of Sciences, St.Petersburg 194021, Russia

A model of chemical bonding between ions in manganites involving covalent one-electron σ bonding is suggested. The covalent one-electron σ bonding gives rise to a strongly correlated state of electrons resulting from the exchange interaction between electrons when they are simultaneously at cation and anion orbitals. The manifestation of the correlated state is the spin and spatial ordering of the electrons resulting in the formation of a spin-ordered electron lattice. The conductivity of manganites in this model is the consequence of displacement of the electron lattice (or its part) from one localization site to another and depends on the type of spin ordering of the electrons in the electron lattice and the localization energy determined by the energy of the one-electron σ bond. The model also assumes a strong polarization of an anion by cations, which facilitates the 3s2p hybridization of the anion and transition of one of the pairs of 2p electrons from the singlet state into the triplet state. This transition leads to formation of the spin-polarized electron lattice (electron spins are parallel) and ferromagnetic ordering of manganese ions. In the model, the effect of colossal magnetoresistance is explained by a change of the conductivity mechanism on application of an external magnetic field, i.e., transition from the conductivity mechanism typical of an ionic crystal to the conductivity provided by the spin-polarized electron lattice.

I. INTRODUCTION

Electric and magnetic properties of doped manganites have received much attention of researchers in recent years because these compounds are thought to be promising for the use as high-sensitivity detectors of magnetic fields, in magnetic resistive memory, readout magnetic heads, and in devices based on the spin-polarized transport.¹

Of special interest among doped manganites are the compounds where transition to the magnetic ordering is accompanied by a metal-insulator transition.² The interplay of magnetic and transport properties in manganites has been explained for a long time in the framework of the double-exchange theory³ which assumes electron exchange between neighboring Mn^{3+} and Mn^{4+} ions via an intervening O^{2-} ion.

However, experimental data obtained in recent years, such as an increase in T_c under an external magnetic field⁴, the effect of colossal magnetoresistance⁵, coexistence of charge ordering and ferromagnetism⁶, ferromagnetic ordering in the compounds $CaCu_3MnO_{12}$ ⁷ and $Tl_2Mn_2O_7$ ⁸ where there are no Mn^{3+} ions, have thrown doubt on the concept of double exchange as a basic mechanism of ferromagnetic ordering and conductivity in manganites⁹.

An important achievement in studies of doped manganites was detection of local distortions of the crystal lattice below T_c .¹⁰ It was shown that the actual crystal lattice strongly differs from the averaged lattice identified by X-ray diffraction. These experimental data have led to the conclusion on a close relation between magnetic and structural properties. However, the questions of how the local lattice distortions influence the electron structure and magnetic ordering and what is the origin of these distortions still remain unresolved.

The goal of this work was to show that the interplay between magnetic, transport, and structural properties can be explained if we take into account a strong covalency of bonding between manganese and oxygen ions and represent it as formation of one-electron σ bonds between these ions. This concept of one-electron σ bonding in manganites was earlier invoked to explain antiferromagnetic ordering.¹¹ In the paper it is shown that this concept can prove useful for explanation of ferromagnetic ordering as well. Moreover, this work has a more general purpose - to show by using manganites as an example how a strongly correlated state of the electrons responsible for magnetic ordering and high conductivity can arise via the exchange interaction between the electrons forming one-electron σ bonds.

II. COVALENT ONE-ELECTRON BONDING

As for the type of chemical bonding, manganites occupy an intermediate position between ionic and covalent crystals. This is evidenced by the difference between electronegativities (1.55 for Mn and 3.44 for O according to Pauling¹²) equal to 1.9, which corresponds to approximately 40% of covalency and 60% of ionicity of bonding between Mn and O atoms. A strong covalency in manganites has been recently experimentally confirmed by high-resolution Mn k_β emission spectroscopy.¹³ Such covalency must manifest itself first of all in the behavior of valence electrons because covalency means sharing of electrons by adjacent ions, while ionicity requires that they be localized at anions.

The question arises how the state of valence electrons in a solid when they are simultaneously localized at anions and shared with cations can be imagined.

For molecules, this type of bonding was regarded as a resonance state of structures with a purely covalent and purely ionic bond.¹² However, it is hardly probable that this approach is appropriate for solids because it is unlikely that a typical two-electron σ bond considered in the resonance model is possible in an ionic crystal. In addition, it is difficult to represent ordering of different types of bonds. Probably, it would be more reasonable to treat such bonding as *sharing of only one electron* by each cation-anion pair.

Let us suppose that at high temperatures, ions are mainly coupled ionically, and this ionic bonding determines the averaged crystal structure of manganites with a high-symmetry lattice of ions. As temperature decreases, additional covalent one-electron σ bonding between Mn and O ions appears. In this case the crystal lattice undergoes distortions, and the crystal symmetry is lowered. Appearance of covalent one-electron bonding between ions means a partial substitution of ionic bonding by covalent bonding, i.e., the Coulomb interaction energy between ions is partly substituted by the covalent bonding energy. As a result, the bonding energy increases, which is evidenced by a decrease in the distance between ions which turns to be less than the sum of the ionic radii. Let us treat this covalent one-electron bonding as a *set of localized bonds* between neighboring ions.

If Mn ions are surrounded by oxygen octahedra, they can use six equivalent hybrid $3d^24s4p^3$ orbitals for covalent bonding. Since the energy level of these hybrid orbitals is higher than the level of d_{xy} , d_{yz} , and d_{xz} orbitals, which results from both admixing of a higher-energy 4p state and splitting of d levels in the octahedral crystal field, it can be supposed that the difference between the energy levels of hybrid and unmixed orbitals is sufficiently large to make the transformation of Mn^{3+} ion from the high-spin state ($t_{2g}^3e_g^1$) into the *low-spin state* (t_{2g}^4) energetically favorable. As a result, all six hybrid orbitals of Mn^{3+} ions will be empty and can be available for covalent bonding. The probability of the low-spin state for Mn^{3+} ion in manganites was also discussed in Ref.¹⁴.

An O^{2-} ion in the manganite structure is also octahedrally surrounded, two of the surrounding cations being Mn ions lying along one of the axes of the octahedron. To provide one-electron σ bonding with these cations, the O^{2-} ion can use its 2p orbital or *two hybrid 3s2p orbitals* pointed to Mn ions. The use of the *2p orbital* has already been discussed¹¹, but the possibility of 3s2p hybridization needs an explanation. Due to a large difference between energies of 2p and 3s levels in an oxygen *atom*, the 3s2p hybridization is thought to be impossible. However, for the O^{2-} ion that is in the *ionic crystal lattice*, there are some factors that facilitate such hybridization. First, the energy level of the 2p state in the O^{2-} ion is higher than that in the oxygen atom, which is evidenced by a difference between the ionic oxygen radius and its atomic radius (1.26Å and 0.9Å, respectively). Second, O^{2-} ion located between two manganese ions with a high positive charge and relatively small ionic radius will be strongly *polarized* by these two cations. The energy of this polarization can partly compensate for the energy spent on excitation of two 2p electrons into the 3s2p state. At last, the energy spent on hybridization will also be compensated by the total gain in energy due to formation of a stronger bonding between ions.

Let us compare covalent one-electron bonds involving 2p and 3s2p orbitals of the O^{2-} ions. We designate them as $1e - \sigma$ and $1e - \sigma^*$, respectively. Since the $1e - \sigma$ bond is formed by a shorter 2p orbital, its energy will be higher than the energy of the $1e - \sigma^*$ bond, and therefore the $1e - \sigma$ bond is more favorable for the crystal stability. However, the $1e - \sigma$ bond can be formed only if a cation and an anion can closely approach each other to provide overlapping of the cation orbital with the 2p anion orbital. The possibility of such overlapping depends on the deformability of the ionic lattice and also on the filling of the cation electron shell. *Nonsphericity* of the electron shell of a Mn^{3+} ion in the low-spin state resulting from the presence of a t_{2g} electron pair in one of the planes hinders a close approach of this cation and anion in this plane because of a strong repulsion between the electron pair and the anion. For this reason, in this plane, only overlapping with a more extended 3s2p orbital and, hence, formation of only $1e - \sigma^*$ bond is possible.

Another difference between $1e - \sigma$ and $1e - \sigma^*$ bonds is the spin state of the anion electron pair that forms these bonds. For $1e - \sigma$ bonds, the pair is in the *singlet* state because it belongs to one p orbital (Pauli exclusion principle); and for $1e - \sigma^*$ bonds the pair is in the *triplet* state because in hybridization two electrons of one 2p orbital must pass into two hybrid 3s2p orbitals, and their spins must be parallel according to the Hund's rule because these orbitals are orthogonal and share a space.

III. SPIN ORDERING OF ELECTRONS AND ELECTRON CRYSTALLIZATION

If two one-electron bonds are formed by one anion with two neighboring cations, the spin state of the anion electron pair participating in covalent bonding will determine the orientation of electron spins in the hybrid orbitals of these cations (Fig. 1).




FIG. 1. Formation of covalent one-electron bonding between Mn^{3+} and O^{2-} ions: (a) $1e - \sigma$ and (b) $1e - \sigma^*$. Empty d^2sp^3 hybrid orbitals of two Mn^{3+} ions and the p orbital occupied by a singlet electron pair of O^{2-} ion (a) and empty d^2sp^3 hybrid orbitals of two Mn^{3+} ions and two $3s2p$ hybrid orbitals occupied by a triplet pair of O^{2-} ion (b) are shown.

Each cation which has six hybrid orbitals can share six electrons - one electron from each of six neighboring anions. The spins of all these six electrons must be parallel according to the Hund's rule for hybrid d^2sp^3 cation orbitals. Therefore, if all the anions and all Mn ions are coupled by one-electron σ bonds, all the electrons taking part in formation of these bonds will be spin-ordered (Fig. 2).




FIG. 2. Electron spin ordering in hybrid orbitals of Mn^{3+} (black circles) and Mn^{4+} (a black square) ions at formation of $1e - \sigma$ and $1e - \sigma^*$ bonds via intervening O^{2-} ions (singlet and triplet pairs). For simplicity, only four of six hybrid orbitals for each cation and only three anions are shown.

Since the Hund's rule reflects the dependence of the electron system energy on the mutual spin orientation, and such dependence is equivalent to the existence of an additional interaction between electrons which is known as the exchange interaction, it can be stated that spin ordering of the electrons participating in covalent bonding results from two exchange interactions - between electrons in the anion orbital (or two hybrid orbitals) and between electrons in the cation hybrid orbitals. *In the case of one-electron bonding*, the sign of the exchange interaction in the cation hybrid orbitals is *always positive*, while the sign of the exchange interaction between the electrons in the anion orbitals, as shown above, can be *positive* (a triplet pair) or *negative* (a singlet pair) (Fig. 2).

In the case of positive signs of the exchange interactions between electrons in both the anion and cation orbitals, the spins of all the electrons participating in one-electron bonding will be parallel, i.e., there will be 100-% spin polarization of the electrons. In the case of different signs of the exchange interactions in cation and anion orbitals, the electrons are also spin-ordered, but their spins are antiparallel in the hybrid orbitals of neighboring cations.

The exchange interaction leads to not only spin ordering of electrons, but also spatial ordering of electrons relative to each other, which can probably be regarded as the electron lattice formation (electron crystallization). On the whole, such ordering resembles the Wigner crystal formation¹⁵, but there is a significant difference. The Wigner crystal

involves a purely Coulomb repulsion between electrons in a uniform field of positive charge. As a result, the Wigner crystal is possible only at low temperatures when the Coulomb repulsion energy exceeds the electron thermal energy, and at these temperatures the crystal acquires the ability to displace as one unit (in the absence of the inhomogeneities that cause pinning). In the case of formation of the electron lattice considered in this paper, the repulsion energy is a *sum* of the purely Coulomb repulsion energy and the repulsion energy due to the exchange interaction between electrons with parallel spins in hybrid orbitals. This leads to electron crystallization at such high temperatures as T_c . Moreover, participation of electrons in covalent bonding causes localization of the electron lattice. The energy of this localization depends on the one-electron bond energy. Hence, the electrons participating in one-electron bonding between ions are in a *strongly correlated state* - they form a *localized spin-ordered electron lattice*.

It is important to mention here that magnetic ordering of inherent magnetic moments of cations resulting from unpaired t_{2g} electrons will be established via a purely magnetic interaction between these magnetic moments and the magnetic moment of the spin-ordered electron lattice. In the case of its 100-% spin polarization (spins are parallel), magnetic moments of all the cations will be ordered ferromagnetically.

IV. CONDITIONS FOR CONDUCTIVITY

Let us consider now the conductivity in the system of strongly correlated electrons. It is likely that, in the presence of strong repulsion between spin-polarized electrons (there cannot be two electrons with parallel spins in one orbital), the only possible conductivity mechanism is displacement of the electron lattice as one unit, i.e., *collective transport of all the electrons from one localization site to a neighboring site*. However, such collective displacement is possible if transition of any electron to a hybrid orbital of a neighboring cation is not forbidden by the spin direction in this orbital and if the localization energy of the electron lattice is not too high. The first condition can be satisfied if the electron lattice is spin-polarized (spins are parallel). The second condition can be fulfilled by a decrease in the extent of overlapping between cation and anion hybrid orbitals, i.e., by an increase in the distance between $Mn^{3+}-O^{2-}-Mn^{3+}$ ions, for instance, due to local crystal lattice distortions observed in doped manganites. However, these conditions contradict each other because weakening of covalent bonds and their breaking due to thermal ion vibrations can lead to a loss of spin ordering between electrons of neighboring cations since spin ordering is created and maintained just by the covalent bonding. Nevertheless, these two conditions can be satisfied simultaneously if the functions of conductivity and maintaining spin ordering are fulfilled by different parts of the spin-polarized lattice. This can be realized if a part of the spin-polarized electron lattice (spins are parallel) remains three-dimensional and localized, while the second part has a lower localization energy and is characterized by a lower dimensionality. The first part of the lattice will provide spin ordering, and the second part will provide conductivity. Below we show how such a separation of functions between the parts of a spin-polarized electron lattice is achieved in doped manganites.

V. MAGNETIC ORDERING AND CONDUCTIVITY IN DOPED AND UNDOPED MANGANITES

Let us now see how magnetic ordering and conductivity of doped and undoped manganites can be explained in the framework of the suggested model.

An experimentally observed perovskite cubic lattice of $CaMnO_3$ with a distance between ions less than the sum of ionic radii (1.865Å and 1.93Å, respectively) and antiferromagnetic ordering of Mn^{4+} ions below T_N are in good agreement with the assumption of six equivalent $1e-\sigma$ bonds formed by each Mn^{4+} ion with six surrounding anions via electrons of singlet pairs. The electrons participating in covalent bonding and being in hybrid cation orbitals prove to be mutually ordered, and their state can also be regarded as an electron lattice. The electron lattice in this case consists of electron octahedra surrounding Mn^{4+} ions, the electron spins being parallel in each octahedron and antiparallel with respect to a neighboring octahedron below T_N . Conductivity is impossible because of different spin orientations of electrons in the orbitals of neighboring cations, consistent with experimental data.²

As to $LaMnO_3$, the nonsphericity of the electron shell of Mn^{3+} ion, as discussed above, results in formation of four $1e-\sigma^*$ bonds in the plane where a pair of t_{2g} electrons is located, which must lead to ferromagnetic ordering between neighboring cations. In the direction normal to this plane, two $1e-\sigma$ bonds can be formed, and their formation will result in antiferromagnetic ordering between neighboring cations. The orientational ordering of Mn^{3+} ions in one plane provides ferromagnetic ordering of all Mn^{3+} ions in this plane, the magnetic ordering between such planes being antiferromagnetic. Precisely this magnetic ordering was observed experimentally below T_N .² The orientational ordering of Mn^{3+} ions in one plane decreases the deformational energy of the ionic lattice when bonds of different lengths ($1e-\sigma^*$ and $1e-\sigma$) are formed. Such ordering is energetically favorable.

The spin-ordered electron lattice in this case consists of electron octahedra surrounding Mn^{3+} ions. The electron spins are parallel in all the electron octahedra lying in the plane of the orientational ordering of Mn^{3+} ions and are antiparallel to the spins of the electron octahedra in the neighboring planes. $LaMnO_3$ is an insulator², and this is likely to be due to both a high localization energy of the electron lattice in the absence of local lattice distortions and a high dimensionality of the spin-polarized lattice (two-dimensional).

In doped manganites which contain both ions (Mn^{3+} and Mn^{4+}), each ion will tend to form the bonds similar to those in undoped compounds. As to coupling between Mn^{3+} and Mn^{4+} ions via an intervening anion, the $1e - \sigma$ bond is likely to be formed between them because it is energetically favorable, and both ions are capable of forming such bond. Such bonding leads to antiferromagnetic ordering of Mn^{3+} and Mn^{4+} ions. Since a Mn^{3+} ion cannot form all the six $1e - \sigma$ bonds, this ion will be coupled with the neighboring Mn^{3+} ions by $1e - \sigma^*$ bonds to result in ferromagnetic ordering between these ions. Between Mn^{4+} ions, $1e - \sigma$ bonds can be formed if deformation of the ionic lattice (or local distortions) is possible, or $1e - \sigma^*$ bonds can be formed if such deformation is impossible.

In this respect, the $A_{0.67}B_{0.33}MnO_3$ chemical composition of doped manganites is unique because it makes possible such charge ordering of Mn^{3+} and Mn^{4+} ions in which each Mn^{4+} ion is surrounded by six Mn^{3+} ions, and each Mn^{3+} ion is surrounded by three Mn^{4+} ions and three Mn^{3+} ions. In the latter case the cations with equivalent oxidation states are the nearest neighbors (Fig. 3).




FIG. 3. Coordination polyhedron of a Mn^{3+} ion in $A_{0.67}B_{0.33}MnO_3$. Three pairs of $1e - \sigma$ bonds with neighboring Mn^{4+} ions and three pairs of $1e - \sigma^*$ bonds with neighboring Mn^{3+} ions via an intervening O^{2-} ions are shown.

Such charge ordering is accompanied by formation of $1e - \sigma^*$ bonds between all Mn^{3+} ions via intervening O^{2-}

ions and $1e - \sigma$ bonds between all Mn^{3+} and Mn^{4+} ions. Mn^{4+} ions are not the nearest neighbors to each other. In Fig. 4, two sublattices formed by Mn^{3+} and Mn^{4+} ions separately due to their charge ordering can be seen. The electron lattice involves in this case two electron sublattices with antiparallel spins.




FIG. 4. Ordering of Mn^{3+} ions (black circles), Mn^{4+} ions (black squares), $1e - \sigma$ bonds formed by singlet pairs (dashed lines), and $1e - \sigma^*$ bonds formed by triplet pairs (solid lines) in $A_{0.67}B_{0.33}MnO_3$. Dashed open circles show O^{2-} ions in the $3s2p$ hybrid state. Local distortions are not shown.

One sublattice consists of electron octahedra around Mn^{4+} ions, and the other electron sublattice comprises electron octahedra around Mn^{3+} ions. Such classification of the electron sublattices is convenient for determining the type of magnetic ordering of intrinsic magnetic moments of cations. In this case Mn^{3+} ions themselves and also Mn^{4+} ions themselves are ordered ferromagnetically, and antiferromagnetically with respect to each other.

On the other hand, to *explain conductivity*, it would be useful to divide the spin-ordered electron lattice into two sublattices formed by electrons of triplet pairs and singlet pairs of anions separately because there is a difference between the localization energies of the electrons of these pairs due to the different energies of $1e - \sigma^*$ and $1e - \sigma$ bonds. Such division of the electron lattice allows one to separate out the less localized part of the spin-polarized electron lattice (spins are parallel). With such choice of electron sublattices, it should be borne in mind that the spins of all the electrons in the sublattice of singlet pairs are ordered. As can be seen from Figs. 3 and 4, the electron sublattice of triplet pairs is a part of the spin-polarized lattice consisting of electron octahedra around Mn^{3+} and turns out to be quasi-one-dimensional in each MnO_2 plane. It is arranged between the nearest Mn^{3+} ions in the form of stripes parallel to each other. These stripes are separated from each other in each MnO_2 plane by the stripes of the electron sublattice of singlet pairs. Since formation of short $1e - \sigma$ bonds between Mn^{3+} and Mn^{4+} ions results

in an increase in the distance between Mn^{3+} ions (local distortions, see Fig. 5), which is accompanied by the decrease in overlapping and hence in the energy of the $1e - \sigma^*$ bond, it can be supposed that in this manganite composition the electron sublattice of triplet pairs becomes less localized and more capable of displacing from one localization site to a neighboring one compared with $LaMnO_3$.




FIG. 5. Projection of a distorted crystal lattice onto a MO_2 plane. The $1e - \sigma$ bonds and $1e - \sigma^*$ bonds formed by singlet and triplet electron pairs are shown. The O^{2-} ions forming the bonds are not shown. The designations are the same as in Fig. 4.

The electron sublattice of singlet pairs remains insulating and preserves spin ordering because a half of electrons of this sublattice is in the hybrid orbitals of the same Mn^{3+} ions in which the electrons of triplet pairs are present (Fig. 3).

VI. FERROMAGNETIC FLUCTUATIONS AND THE EFFECT OF EXTERNAL MAGNETIC FIELD ON T_c

The above discussion shows that T_c is the temperature of formation of a spin-ordered electron lattice incorporating a spin-polarized electron sublattice of triplet pairs. It is evident that this temperature must be preceded by a temperature interval in which electrons are still localized in anion orbitals. Though anions polarized by Mn ions can be in the $3s2p$ hybridization state, static overlap of these hybrid orbitals with hybrid cation orbitals is absent. Only dynamic overlap due to thermal vibrations of ions is possible. It can manifest itself as short-range ferromagnetic fluctuations observed experimentally.¹⁶ In this temperature interval, the "foundation" for formation of static $1e - \sigma^*$ bonds and a spin-polarized sublattice via a gradual spin polarization of all the electrons of triplet pairs is laid. Apparently, application of an external magnetic field, which aligns spins of all unpaired electrons, can accelerate the formation of the spin-polarized electron sublattice and thereby increase T_c . For similar reasons, T_c must also increase with increasing magnetic field.¹⁷

In terms of the suggested model, the effect of colossal magnetoresistance is the consequence of the change of the

conductivity mechanisms, i.e., from the thermally activated conductivity typical of a solid with preferentially ionic bonding to the metallic conductivity caused by a spin-polarized electron sublattice of triplet pairs whose formation is facilitated by the external magnetic field.

VII. CONCLUSIONS

The above consideration shows that the origin of a strongly correlated state of electrons in manganites responsible for magnetic ordering and metallic conductivity can be understood if we represent a strong covalency of chemical bonding between manganese and oxygen ions by formation of one-electron σ bonds between these ions. These bonds are formed by overlapping of empty d^2sp^3 orbitals of Mn ions with orbitals of anions. Each anion can form two one-electron σ bonds with two neighboring Mn ions. It is assumed in this case that a Mn^{3+} ion prefers a low-spin state and can polarize an anion thereby causing its 3s2p hybridization which is accompanied by transition of its singlet electron pair into a triplet state. Both singlet and triplet pairs can participate in covalent one-electron bonding, which results in different spin ordering of electrons in hybrid orbitals of neighboring cations. The correlated state of the electrons forming one-electron σ bonds arises due to the exchange interaction between electrons when they are simultaneously in hybrid orbitals of a cation and in the p orbitals (or hybrid 3s2p orbitals) of anions. This exchange interaction leads to both the spin ordering and spatial ordering of electrons, which can be represented as formation of a spin-ordered electron lattice. The conductivity produced by such an electron lattice depends on the type of spin ordering, lattice dimensionality, and the localization energy, which in turn depends on the one-electron σ bond energy.

The model of chemical bonding suggested here can prove useful for explanation of high- T_c superconductivity in cuprates. In¹⁸, an attempt was made to represent the bonding between Cu and O ions whose covalency is stronger than in manganites through a resonance state of ionic and ordinary two-electron σ bonds (in analogy with bonding in molecules¹²). In spite of the fact that such representation led to the model of additional π bonding between Cu and O ions which qualitatively explained high- T_c superconductivity, many assumptions of this model remained unclear and even controversial.

The use of the notion of one-electron σ bonding to explain superconductivity can also clarify the picture of the behavior of correlated electrons in cuprate superconductors. With this notion, it becomes clear that, in the CuO_2 plane, formation of one-electron σ bonds is also possible. The anion is also subjected to polarization by cations. However, for superconductivity, the anion polarization in a strongly asymmetric crystal field normal to the CuO_2 plane proves to play a decisive role. It promotes the 3s2p hybridization of an anion and forces its triplet electron pair to transfer from two hybrid orbitals into one hybrid orbital pointed to a positively charged plane (for instance, Y^{3+}). In this case, the spin state of this electron pair is transformed from a triplet state into a singlet one. These singlet pairs which are in the 3s2p hybrid orbitals of anions form a localized electron lattice. Displacement of this lattice, i.e., collective displacement of singlet pairs, can occur after π overlapping of these 3s2p hybrid orbitals with empty hybrid orbitals of Cu^{3+} ions and formation of delocalized π orbitals. Thus the pairing mechanism in high- T_c superconductors can be the exchange interaction between electrons in a hybrid orbital of an anion polarized by a strongly asymmetric crystal field. In this model, the mechanism of superconductivity is displacement of the lattice of electron pairs in delocalized π orbitals that belong to the chains of Cu^{3+} and O^{2-} ions in the CuO_2 plane.

* Corresponding author: M.V.Krasinkova, E-mail: marina.shuv@pop.ioffe.rssi.ru

¹ G. Prinz and K. Hathaway, Phys. Today **48**, 24 (1995). J.L. Simonds, Phys. Today, **48**, 26 (1995). G.Prinz, Phys. Today **48**, 58 (1995).

² G.H. Jonker and J.H. Van Santen, Physica (Amsterdam) **16**, 337 (1950). E.O. Wollen and W.C. Koehler, Phys.Rev. **100**, 548 (1955). R.H. Heffner, L.P. Le, M.F. Hundley, and J.J. Neumeier, G.M. Luke, K. Kojima, B. Nachumi, and Y.J. Uemura, D.E. MacLaughlin, S.-W. Cheong, Phys.Rev.Lett. **77**, 1869 (1996).

³ C. Zener, Phys.Rev. **82**, 403 (1951). P.W. Anderson and H. Hasegawa, Phys.Rev. **100**, 675 (1955).

⁴ J. Fontcuberta, B. Martinez, A. Seffar, S. Pinol, J.L. Garcia-Munoz, and X. Obradors, Phys.Rev.Lett. **76**, 1122 (1996).

⁵ R. von Helmolt, J. Wecker, B. Holzapfel, L. Schultz, and K. Samwer, Phys.Rev.Lett. **71**, 2331 (1993). S. Jin, T.H. Tiefel, M. McCormack, R.A. Fastnacht, R. Ramesh, and L.H. Chen, Science **264**, 413 (1994). K. Chahera, T. Ohno, M. Masahiro, and Y. Kozono, Appl.Phys.Lett. **63**, 1990 (1993).

⁶ C.H. Chen and S.-W. Cheong, Phys.Rev.Lett. **76**, 4042 (1996). P.G. Radaelli, D.E. Cox, L. Capogno, S.-W. Cheong, and M. Marezio, Phys.Rev. B **59**, 14 440 (1999).

- ⁷ Z. Zeng and M. Greenblatt, M.A. Subramanian, M. Croft, Phys.Rev.Lett. **82**, 3164 (1999).
- ⁸ Y. Shimakawa, Y. Kubo, and T. Manako, Nature **379**, 53 (1996).
- ⁹ A.J. Millis, P.B. Littlewood, and B.I. Shraiman, Phys.Rev.Lett. **74**, 5144 (1995).
- ¹⁰ D. Louca and T. Egami, E.L. Brosha, H. Roder, and A.R. Bishop, Phys.Rev. B **56**, R8475 (1997). S.J.L. Billinge, R.G. DiFrancesco, G.H. Kwei, J.J. Neumeier, and J.D. Thompson, Phys.Rev.Lett. **77**, 715 (1996). D. Louca and T. Egami, Phys.Rev. B **59**, 6193 (1999).
- ¹¹ J.B. Goodenough, Phys.Rev. **100**, 564 (1955).
- ¹² L. Pauling, *The Nature of the Chemical Bond and the Structure of Molecules and Crystals*, 2nd ed. (Cornell Univ. Press, Ithaca, New York, 1940).
- ¹³ T.A. Tyson and Q. Qian, C.-C. Kao, and J.-P. Rueff, F.M.F. de Groot, M. Croft, S.-W. Cheong, M. Greenblatt, M.A. Subramanian, Phys.Rev. B **60**, 4665 (1999).
- ¹⁴ T.H. Geballe and B.Y. Mozyzhes, J.Electroceramics **4**, 289 (2000).
- ¹⁵ E. Wigner, Phys.Rev. **46**, 1002 (1934).
- ¹⁶ J.W. Lynn, R.W. Erwin, J.A. Borchers, G. Huang, A. Santoro, J.I. Peng, and Z.Y. Li, Phys.Rev.Lett. **76**, 4046 (1996).
- ¹⁷ A. Urushibara, Y. Moritomo, T. Arima, A. Asamitsu, G. Kido, Y. Tokura, Phys.Rev. B **51**, 14 103 (1995).
- ¹⁸ M.V. Krasinkova, Pis'ma Zh. Tekh. Fiz. **23**, 57 (1997) [Sov.Tech.Phys.Lett. **23**, 681 (1997)]. M.V. Krasinkova, Zh. Tekh. Fiz. **68**, 82 (1998) [Sov.Phys.Tech.Phys. **43**, 1347 (1998)].